

Parameterized optimized effective potential for atoms

A. Sarsa, F. J. Gálvez and E. Buendía
*Departamento de Física Moderna, Facultad de Ciencias,
Universidad de Granada, E-18071 Granada, Spain*
(Dated: February 2, 2008)

The optimized effective potential equations for atoms have been solved by parameterizing the potential. The expansion is tailored to fulfill the known asymptotic behavior of the effective potential at both short and long distances. Both single configuration and multi configuration trial wave functions are implemented. Applications to several atomic systems are presented improving previous works. The results here obtained are very close to those calculated in either the Hartree-Fock and the multi configurational Hartree-Fock framework.

PACS numbers: 31.10.+z, 31.15.-p, 31.15.Pf, 31.15.Ne, 02.70.-c

I. INTRODUCTION

The Hartree-Fock (HF) method is the best approximation to the atomic and molecular problem in the independent particle model with a single configuration. Originally it was formulated in terms of a single Slater determinant but in current applications a single configuration, with one or more Slater determinants coupled in the LS scheme, is used, see e.g. [1]. An alternative to the HF equations, always within the independent particle model, is given by the Optimized Effective Potential (OEP) method proposed by Sharp and Horton [2] as a variational alternative to the Slater's simplified treatment of the exchange term in the HF equations based on the averaging in the occupied orbitals [3].

In the OEP method an additional constraint is imposed into the variational problem: the orbitals must satisfy a single-particle Schrödinger equation with a certain local potential, the same for all the electrons (with the same spin). The expectation value of the hamiltonian of the N -electron system becomes a functional of such a local potential. The effective potential is then varied to minimize the total energy. This gives rise to a linear integral equation in the effective potential [2, 4] whose solution gives the optimized effective potential. The wave function of the system is then constructed from the single-particle wave functions which are eigenfunctions of the optimized effective potential.

The OEP method has been used in connection with the Kohn-Sham density functional theory [5, 6, 7, 8]. Accurate spin-polarized exchange-only Kohn-Shan potential has been constructed with the OEP method [9]. The local exchange potential obtained from the OEP method has many of the analytical features of the exact Kohn-Sham potential, and it has been recognized as the exact implementation of exchange-only density functional theory [10]. For example the HF potential can support very few unoccupied excited states because of its exponential fall off and therefore it is not a good starting point to describe excited states. On the other hand as the OEP potentials presents the proper long range behavior it provides better excitation energies.

The use and application of the original form of the

OEP was hindered because of the computational difficulties posed by the OEP integral equation. Initially, a numerical grid method was employed by Talman and Shadwick [4] to solve this equation and further more accurate calculations for atoms have been carried out by using a more refined mesh [11, 12]. Also some approximations to the OEP equations were proposed [13]. Very recently an iterative scheme based on solving the partial differential equations satisfied by the orbitals shift that satisfy exactly the Krieger, Li and Iafrate approximation has been devised [14]. An alternative methodology to board this problem uses a finite basis set expansion and/or analytical forms for the effective potential [8, 15, 16, 17, 18, 19, 20, 21, 22]. The parameterization of the effective potential simplifies greatly the numerical problem of solving the integral equations [8, 16, 19, 22]. The explicit form or the parameterization is proposed by using different arguments [16, 19, 23]. The parameters can be fixed either by using the variational method or by matching the one-body eigenvalues of the corresponding spin-averaged Dirac equation to the experimentally observed one-electron ionization energies [17].

The aim of this work is to obtain the energy and other properties for atoms in the parameterized OEP approximation. The analytical form used for the optimized potential has been used previously [15, 17] but with simpler expansions. In this work we increase the number of free parameters in the effective potential until convergence is reached. In this way we improve previous results obtained not only within the parameterized OEP scheme but also by solving numerically the integral equation. We have studied the ground state of the atoms Li to Ar in the LS coupling. Finally the OEP approximation is extended here to multi configuration wave functions in the same spirit of the Multi Configuration Hartree Fock method. The binding energy, single particle energies and exchange energies are obtained and compared systematically with their Hartree-Fock counterparts. The best results for the energy of the present work are roughly 1 mhartree above the best self consistent field results showing the good performance of the local potential approximation.

The structure of this work is as follows. In Section II we show in detail the parameterization and the algorithm

used. The results obtained are reported and discussed in Section III. The conclusions and perspectives of this work can be found in Section IV. Atomic units are used throughout.

II. PARAMETERIZED OPTIMIZED EFFECTIVE POTENTIAL

The single-particle wave functions used to construct the Slater determinants of a given single configuration are the eigenfunctions of the so called effective potential, that in this work is taken to be central,

$$\left(-\frac{1}{2}\vec{\nabla}^2 + V_e(r)\right) \phi_{\lambda,\sigma}(\vec{r}) = \epsilon_\lambda \phi_{\lambda,\sigma}(\vec{r}) \quad (1)$$

where λ and σ stand for the spatial and spin quantum numbers, respectively.

The total energy of the system is therefore a functional of this single particle potential. The minimum condition on this potential leads to the OEP equations [2] fixing the best effective potential. The energy so obtained will be an upper bound to the exact one and it will be above the HF value. It is also worth pointing out that the role of the effective potential here is just an auxiliary function used to calculate the orbitals in the wave function.

The parameterization chosen in this work for the effective potential is

$$V_e(r) = -\frac{1}{r} \left(Z - N + 1 + (N-1) \sum_{p=0}^S \sum_{k=1}^{n_p} c_{k,p} r^p e^{-\beta_{k,p} r} \right) \quad (2)$$

with the condition

$$\sum_{k=1}^{n_0} c_{k,0} = 1 \quad (3)$$

imposed in order to match the correct short-range behavior of the potential. This constraint makes that we must deal with, at least, two terms with $p = 0$, i.e. $n_0 \geq 2$. With respect to the long range asymptotic behavior, this functional form is such that the potential goes as $-(Z - N + 1)/r$ for large electron-nucleus distances. The number of basis functions used to expand the effective potential is incremented systematically until convergence is reached. As we shall see later the importance of considering higher values of p increases with the number of electrons. Finally it is worth mentioning here that regular behavior close to the nucleus has shown to be relevant in local effective potential theories [24].

The algorithm is the following

1. Pick the initial values of the effective potential parameters, $c_{k,p}$ and $\beta_{k,p}$ such that the short and long range asymptotic behaviors are satisfied.
2. Solve the one-electron Schrödinger equation for each occupied orbital.

3. Use these orbitals to build up the N-electron wave function and calculate the expectation value of the Hamiltonian.
4. Optimize the total energy with respect to the free parameters ($c_{k,p}$, $\beta_{k,p}$) with the constraint given by equation 3.
5. Increase the size of the expansion of the effective potential until convergence.

The single-particle Schrödinger equation 1 in step 3 is solved by expanding the radial orbitals in a basis set of Slater type functions as in the Roothaan-Hartree-Fock method. In particular we have used the same size of the basis set as in reference [25]: the s -type single particle orbitals are expanded by using 6 basis functions for the Li to Ne atoms and 8 basis functions for the Na to Ar atoms. The radial part of p -type single particle orbitals is developed as the sum of 4 basis functions for the B to Ne atoms, 5 basis functions for Na and Mg and 8 basis functions for the atoms Al to Ar. By doing this all the different matrix elements can be computed analytically. The total energy has been minimized with respect to the free parameters of the effective potential by using a SIMPLEX [26] algorithm.

Once obtained the single particle wave functions we can calculate some other quantities of interest such as the exchange energy

$$E_x = -\frac{1}{2} \sum_{\sigma} \sum_{\lambda,\mu=1}^{N_{\sigma}} \int d\vec{r} \int d\vec{r}' \frac{\phi_{\lambda\sigma}^*(\vec{r}) \phi_{\mu\sigma}^*(\vec{r}') \phi_{\mu\sigma}(\vec{r}) \phi_{\lambda\sigma}(\vec{r}')}{|\vec{r} - \vec{r}'|} \quad (4)$$

and the Hartree-Fock single-particle energies, $\epsilon_{nl}^{\text{HF}}$, obtained from the expectation value

$$\epsilon_{\lambda}^{\text{HF}} = \mathcal{I}_{\lambda} + \sum_{\mu} (\mathcal{J}_{\lambda\mu} - \mathcal{K}_{\lambda\mu}) \quad (5)$$

where $\mathcal{I}$, $\mathcal{J}$ and $\mathcal{K}$ are the usual single particle, direct and exchange terms calculated starting from the eigenfunctions of the effective potential. The single particle energies $\epsilon_{\lambda}^{\text{HF}}$ of equation 5 does not coincide with the eigenvalues ϵ_{λ} of equation 1, except for the highest occupied orbital. This was proven in reference [27] within a framework of spin-polarized and arbitrary exchange-correlation functionals. In the scheme of the present study this condition states that $\epsilon_{\lambda_h}^{\text{HF}} = \epsilon_{\lambda_h}$, where λ_h stands for the highest occupied level. The fulfillment of this condition has been used previously, e.g. [5, 12] to asses the accuracy in the solution of the OEP equations.

In general the optimization procedure is very stable for both the total energy and the Hartree-Fock values $\epsilon_{\lambda}^{\text{HF}}$. However this is not the case for the eigenvalue ϵ_{λ} , as has been noted previously [9, 12] by using a numerical grid method. For this reason, and because the exact solution of the OEP must also satisfy the virial relation and the exchange-only virial relation [28, 29] we have minimized the quantity

$$\langle H \rangle + |\epsilon_{\lambda_h} - \epsilon_{\lambda_h}^{\text{HF}}| + |E_x - E_x^{\text{vir}}| \quad (6)$$

where E_x^{vr} is given by

$$E_x^{vr} = - \int d\vec{r} \rho(\vec{r}) \vec{r} \cdot \vec{\nabla} V_x(\vec{r}) \quad (7)$$

and V_x is the exchange potential. By doing this no significant changes are found for both the total energy and the Hartree-Fock eigenvalues. One should expect that this procedure gives rise to a better description of the asymptotic region of the optimized effective potential. As it has been previously pointed out [5] these two conditions are in some sense complementary. The main contributions to the quantities involved in the exchange only virial condition arise from the internal region of the atoms whereas the highest energy eigenvalue is governed by the outer region. By including these two conditions in the energy functional we observed a better and more stable convergence in the free parameters as compared to an unconstrained minimization of $\langle H \rangle$. With the basis set used here both conditions are satisfied within one part in 10^{-6} hartree.

The method can be generalized straightforwardly to deal with a Multi Configuration expansion. The starting trial wave function is written as a linear combination of m single configuration wave functions with the total orbital angular momentum and spin of the state under study. The hamiltonian is diagonalized in this set. The orbitals required to build the different Slater determinants are obtained as the eigenfunctions of the single particle effective potential containing several free parameters. The total energy is minimized with respect to those parameters as before. This more general trial wave function will provide not only a better description of the lowest energy state of a given symmetry but also a variational approximation to the excited states because of the Hylleraas-Undheim theorem which states that the eigenvalues constitute upper bounds to the first m bound states. These type of wave functions have been recently used along with a correlation factor of Jastrow type to study some excited states of the beryllium atom and its isoelectronic series [30, 31].

III. RESULTS

In Table I the results obtained by using different parameterizations of the effective potential are analyzed. We show the values for the Ne atom which is representative for the systems studied here. We compare with the numerical optimized effective potential results of Refs. [12] and [32] and with the finite-basis set expansion of the effective potential of Refs. [19, 22]. The approximate values of reference [9] are also reported. The HF results are taken from [33] and are the benchmark values for the total energy obtained from the OEP method. We also report the exchange energy, E_x , and the eigenvalues ϵ_{nl} and $\epsilon_{nl}^{\text{HF}}$. The notation used for the parameterization is $(0^{n_0}, 1^{n_1}, 2^{n_2}, \dots)$ where n_p is the number of functions of the type $e^{-\beta_{k,p} r} r^p / r$ used in the expansion of the effective potential 2.

The best results are obtained with the $(0^2, 1^2, 2^2)$ parameterization, and this is the one used for the rest of the atoms. It can be seen that the value of the total energy does not depend substantially on the basis set, but this is not the case of the other quantities. The use of bigger basis set sizes for the effective potential does not improve noticeably the energy for the atoms Li to Ar and it increases the computational effort. However for heavier atoms we have numerically checked that the rate of convergence can be substantially improved by including higher powers of r in the parameterization of the effective potential.

The best results of this work improve previous ones obtained within the optimized effective potential scheme and it is only 0.57 mhartree above the Hartree-Fock result. The expectation values $\epsilon_{\lambda}^{\text{HF}}$ obtained with the OEP orbitals are in a very good agreement with the corresponding single particle energies obtained within the Hartree-Fock framework. This is because of the fact, pointed out previously [9], that the single particle wave functions calculated from the OEP method are a good approximation to the Hartree-Fock orbitals.

The ground state and exchange energies of the atoms Li to Ar as well as the single-particle expectation values, $\epsilon_{nl}^{\text{HF}}$, and the single-particle OEP eigenvalues, ϵ_{nl} , are reported in tables II and III. All of these quantities are compared with those obtained within the HF framework and with the numerical optimized effective potential results of reference [32]. Within the Hartree-Fock framework the two quantities $\epsilon_{nl}^{\text{HF}}$ and ϵ_{nl} are the same so that we do not list them separately. For all the atoms the $(0^2, 1^2, 2^2)$ parameterization for the effective potential has been used.

As can be noted there is an appreciable improvement of the results, especially for the lightest atoms, which can be more easily seen in Figure 1 where we plot the relative error (in %) of our ground state energy with respect to the HF one as compared to the relative error for the numerical solution. In the present work the relative error is nearly constant for all the atoms considered. In principle one should expect a better energy coming from the numerical solution than from the parametrized one as the used in this work.

Previous works have also reported energies calculated from a parametrized solution that are below the numerical solution [19, 22]. The reason for the better performance of the parametrized solution may be due to numerical inaccuracies in the numerical solution of the OEP equations due to the rather involved procedure of its solution. In an attempt to elucidate this fact we have carried out the following calculation. Starting from the tabulation of Aashamar et al. [32] we have built a parameterized potential of $(0^2, 1^2, 2^2)$ type by fitting the numerical values to that functional form. Then the fitted potential has been used in our code to determine the best orbitals within the effective potential approach of this work without changing the fitted potential. Here we show the results for the Ne atom, which are repre-

TABLE I: Convergence in the parameterization of the effective potential for the Ne atom. The results are compared with the Hartree-Fock values (HF), two different set of results obtained by solving numerically the optimized effective potential (NOEP) and with the Krieger, Li and Iafrate approximation. $(0^{n_0}, 1^{n_1}, 2^{n_2}, \dots)$ stands for n_p functions of the type $e^{-\beta_{k,p} r} r^p / r$ in the expansion.

	HF[33]	NOEP[32]	NOEP[12]	KLI[9]	OEP[19]	OEP[22]
$\langle H \rangle$	-128.54710	-128.5455	-128.5455	-128.5448	-128.5455	-128.5456
E_x	-12.10835	-	-12.1050	-	-12.1068	-12.1055
$\epsilon_{1s}^{\text{HF}}$	-32.77244	-	-	-	-	-
ϵ_{1s}	-	-30.8155	-	-	-	-30.8274
$\epsilon_{2s}^{\text{HF}}$	-1.93039	-	-	-	-	-
ϵ_{2s}	-	-1.71200	-	-	-	-1.7196
$\epsilon_{2p}^{\text{HF}}$	-0.85041	-	-	-0.8494	-0.84975	-
ϵ_{2p}	-	-0.84571	-0.8507	-0.8494	-0.85210	-0.8506
	(0^2)	(0^3)	$(0^2, 1^1)$	$(0^2, 1^1, 2^1)$	$(0^2, 1^2, 2^2)$	
$\langle H \rangle$	-128.54486	-128.54641	-128.54632	-128.54642	-128.54652	
E_x	-12.09600	-12.11006	-12.10020	-12.10002	-12.10591	
$\epsilon_{1s}^{\text{HF}}$	-32.76213	-32.76843	-32.77819	-32.77775	-32.77532	
ϵ_{1s}	-30.98356	-30.77320	-30.83773	-30.90031	-30.87242	
$\epsilon_{2s}^{\text{HF}}$	-1.93007	-1.92862	-1.93405	-1.93386	-1.93102	
ϵ_{2s}	-1.70548	-1.71716	-1.71928	-1.71978	-1.72224	
$\epsilon_{2p}^{\text{HF}}$	-0.85251	-0.84848	-0.85384	-0.85381	-0.85130	
ϵ_{2p}	-0.85251	-0.84848	-0.85384	-0.85381	-0.85130	




Figure 1, Sarsa et al.

FIG. 1: Relative difference (in percent) between the OEP ground state energy of this work and the HF one for the atoms Li through Ar as compared with the same quantity calculated from the numerical results (NOEP) of [32].

sentative of the rest of the atoms analyzed. The total energy obtained in this way is -128.5457 hartree to be compared with -128.5455 hartree reported in Ref. [32]. The difference between these two values is due to a better performance of the parameterized potential which allows one to work analytically. In particular the asymptotic behavior of the potential is taken into account in a more

effective manner.

The difference between the numerical OEP energy and the result obtained here (-128.5465) is more accused. This discrepancy is due to the differences in the effective potential. In Figure 2 we plot the effective potential for the Ne atom and the differences with the numerical results of [32]. The results for Ne are representative for the rest of the atoms considered here. Notice that the differences are multiplied by 10 to better see them in the scale of the figure.




Figure 2, Sarsa et al.

FIG. 2: Effective potential for Ne and the difference (multiplied by 10) with the numerical results of [32].

TABLE II: Energy and single particle energies for He to Ne atoms (OEP). The results are compared with the Hartree-Fock values (HF) obtained from reference [33] and with the numerical solution of the optimized effective potential equation of [32], the exchange energies for the closed shell atoms have been taken from [6].

	Li(² S)	Be(¹ S)	B(² P)	C(³ P)	N(⁴ S)	O(³ P)	F(² P)	Ne(¹ S)
E								
HF	-7.43273	-14.57302	-24.52906	-37.68862	-54.40093	-74.80940	-99.40935	-128.54710
NOEP	-7.4324	-14.5725	-24.5278	-37.6865	-54.3980	-74.8075	-99.4075	-128.5455
OEP	-7.43261	-14.57291	-24.52874	-37.68774	-54.40029	-74.80888	-99.40875	-128.54652
E_x								
HF	-1.78119	-2.66692	-3.74759	-5.04930	-6.59707	-8.18189	-10.01105	-12.10835
NOEP	-	-2.666	-	-	-	-	-	-12.105
OEP	-1.78119	-2.66627	-3.74661	-5.05059	-6.59606	-8.18039	-10.01267	-12.10591
$\epsilon_{1s}^{\text{HF}}$								
HF	-2.47774	-4.73267	-7.69534	-11.32552	-15.62906	-20.66866	-26.38276	-32.77244
OEP	-2.47783	-4.73366	-7.69628	-11.32370	-15.63409	-20.66947	-26.37617	-32.77532
ϵ_{1s}								
NOEP	-2.08186	-4.12785	-6.91330	-10.35324	-14.46742	-19.21792	-24.66731	-30.81549
OEP	-2.09495	-4.06912	-6.92455	-10.42413	-14.61828	-19.30917	-24.78314	-30.87242
$\epsilon_{2s}^{\text{HF}}$								
HF	-0.19632	-0.30927	-0.49471	-0.70563	-0.94532	-1.24432	-1.57254	-1.93039
OEP	-0.19630	-0.30962	-0.49556	-0.70415	-0.94748	-1.24639	-1.56989	-1.93102
ϵ_{2s}								
NOEP	-0.19644	-0.30838	-0.52804	-0.74958	-0.99752	-1.19129	-1.43073	-1.71200
OEP	-0.19630	-0.30962	-0.52968	-0.75608	-1.00990	-1.20004	-1.44285	-1.72224
$\epsilon_{2p}^{\text{HF}}$								
HF	-	-	-0.30986	-0.43334	-0.56759	-0.63191	-0.73002	-0.85041
OEP	-	-	-0.30844	-0.43313	-0.56816	-0.63174	-0.72885	-0.85130
ϵ_{2p}								
NOEP	-	-	-0.30976	-0.43067	-0.56322	-0.62933	-0.72479	-0.84571
OEP	-	-	-0.30844	-0.43313	-0.56816	-0.63174	-0.72885	-0.85130

In Figure 3 we plot, the function $V_p(r)$ obtained from the effective potential as (see Eq. 2)

$$V_p(r) = \frac{rV_e(r) + Z - N + 1}{(N - 1)} \quad (8)$$

We compare the best optimized effective potential obtained in this work with the numerical result calculated from the tabulated values of Ref. [32], the fitted potential is included for the shake of completeness.

Although the effective potential in the numerical approach and the parameterized solution of this work is very similar, see Figure 1, the differences for $V_p(r)$ become apparent. The main discrepancies are for $r \gtrsim 1.5$ au, with a faster decay to zero of the parameterized solution. The magnitude of this part of the effective potential is smaller than the total potential. Therefore, the discrepancies between the parametrized and numerical solutions are not very accused, in this case ~ 0.001 mhartree.

It is worth mentioning here that for Be, Ne, Mg and Ar atoms more accurate numerical results for highest occupied single particle eigenvalues have been reported, [5, 12], but the total energy is very similar to that of reference [32].

To illustrate the performance of the Optimized Effective Potential with a multiconfigurational trial wave func-

tion we have applied it to a simple case. In Table IV we study the ground and the first ¹P excited state of the beryllium atom using a two configuration wave function in both cases. We compare our results with the corresponding Multi Configuration Hartree Fock ones [1, 34] for the same states and configurations. For the ground state, where the 2s-2p near degeneracy plays an important role, we have expanded the wave function in terms of the configurations $1s^22s^2$ and $1s^22p^2$. We have also studied the ¹P state arising from the $1s^22s2p$ configuration by considering the mixing with the $1s^22p3d$ one.

Thus for the ground state, the trial wave function is written as

$$|\Psi\rangle = c_1|1s^22s^2;^1S\rangle + c_2|1s^22p^2;^1S\rangle \quad (9)$$

and for the $1s^22s2p$ ¹P state the wave function is

$$|\Psi\rangle = d_1|1s^22s2p;^1P\rangle + d_2|1s^22p3d;^1P\rangle \quad (10)$$

In Table IV we show the values for the energy and for the coefficients c_k and d_k obtained with the OEP method as compared with the corresponding MCHF ones [34]. It is apparent the good agreement between the two sets of results for both states that illustrates that the OEP method not only provides a good value for the energy but also an adequate weight for any of the configurations involved in the corresponding state. For this case the

TABLE III: The same as in Table II for the atoms Na to Ar.

	Na(² S)	Mg(¹ S)	Al(² P)	Si(³ P)	P(⁴ S)	S(³ P)	Cl(² P)	Ar(¹ S)
E								
HF	-161.85891	-199.61464	-241.87671	-288.85436	-340.71878	-397.50490	-459.48207	-526.81751
NOEP	-161.8565	-199.6115	-241.873	-288.850	-340.714	-397.500	-459.477	-526.810
OEP	-161.85770	-199.61303	-241.87415	-288.85160	-340.71537	-397.50155	-459.47854	-526.81405
E_x								
HF	-14.01752	-15.99429	-18.07225	-20.28350	-22.64091	-25.00614	-27.51653	-30.18494
NOEP	-	-15.988	-	-	-	-	-	-30.175
OEP	-14.01160	-15.98762	-18.06349	-20.27359	-22.62640	-24.99093	-27.49826	-30.17936
$\epsilon_{1s}^{\text{HF}}$								
HF	-40.47850	-49.03174	-58.50103	-68.81246	-79.96971	-92.00445	-104.88442	-118.61035
OEP	-40.47877	-49.03776	-58.50569	-68.81693	-79.97485	-92.01615	-104.89542	-118.60723
ϵ_{1s}								
NOEP	-38.04248	-46.32444	-55.5604	-65.6547	-76.59185	-88.34890	-100.9622	-144.4452
OEP	-37.91728	-46.16018	-55.37384	-65.48096	-76.39372	-88.09753	-100.59527	-114.45720
$\epsilon_{2s}^{\text{HF}}$								
HF	-2.79703	-3.76772	-4.91067	-6.15654	-7.51110	-9.00429	-10.60748	-12.32215
OEP	-2.79384	-3.77025	-4.91136	-6.15701	-7.51443	-9.01030	-10.61285	-12.31768
ϵ_{2s}								
NOEP	-2.48096	-3.09619	-4.15912	-5.34036	-6.63449	-8.01012	-9.51097	-11.14504
OEP	-2.22265	-3.11058	-4.17483	-5.355205	-6.64244	-7.97517	-9.54963	-11.19559
$\epsilon_{2p}^{\text{HF}}$								
HF	-1.51814	-2.28223	-3.21830	-4.25605	-5.40096	-6.68251	-8.07223	-9.57146
OEP	-1.51617	-2.28607	-3.22178	-4.25935	-5.40745	-6.69193	-8.07954	-9.56872
ϵ_{2p}								
NOEP	-1.19797	-1.85896	-2.73146	-3.71956	-4.81814	-5.99544	-7.29541	-8.72568
OEP	-1.16901	-1.86617	-2.74466	-3.73189	-4.82825	-5.96260	-7.33117	-8.76408
$\epsilon_{3s}^{\text{HF}}$								
HF	-0.18210	-0.25305	-0.39342	-0.53984	-0.69642	-0.87953	-1.07291	-1.27735
OEP	-0.18172	-0.25359	-0.38663	-0.54090	-0.69896	-0.88305	-1.07617	-1.27597
ϵ_{3s}								
NOEP	-0.18211	-0.25206	-0.38859	-0.53937	-0.69510	-0.80812	-0.93838	-1.09280
OEP	-0.18172	-0.25359	-0.38388	-0.53904	-0.69884	-0.80924	-0.94641	-1.09886
$\epsilon_{3p}^{\text{HF}}$								
HF	-	-	-0.20995	-0.29711	-0.39171	-0.43737	-0.50640	-0.59102
OEP	-	-	-0.21062	-0.29721	-0.39325	-0.43950	-0.50812	-0.58979
ϵ_{3p}								
NOEP	-	-	-0.20910	-0.29630	-0.38737	-0.43662	-0.50027	-0.58532
OEP	-	-	-0.21062	-0.29721	-0.39325	-0.43950	-0.50812	-0.58979

optimized effective potential method with multi configuration trial wave function compares with the MCHF at the same level as the OEP method with the Hartree-Fock. More complex wave functions using a larger number of configurations have been used to study these and some other excited states of this atom and its isoelectronic series [30, 31]. In those works, not fully optimized multi configuration wave function were used to build up more accurate explicitly correlated wave functions of Jastrow type.

IV. CONCLUSIONS

The optimized effective potential with parameterized potential has been used to study the ground state of the atoms from Li to Ar. Parameterized orbitals have been used to solve the corresponding single particle

TABLE IV: Energy and coefficients for some multi configurational wave functions for the ground and the first excited state of ¹P type for the beryllium atom.

¹ S	E	c_1	c_2
MCHF	-14.61685	0.95003	0.31214
OEP	-14.61637	0.95008	0.31202
¹ P	E	d_1	d_2
MCHF	-14.41156	0.97524	-0.22116
OEP	-14.41131	0.97489	-0.22270

Schrödinger equation. The virial relation of Ghosh and Parr and Levy and Perdew involving the exchange energy and the exchange potential and a condition for the highest energy occupied orbital of Krieger, Li and Iafrate have been imposed. These are two analytically known conditions that the exact solution of the OEP equations must




Figure 3, Sarsa et al.

FIG. 3: $V_p(r)$ defined in Eq. (8) obtained from the best parameterized effective potential of this work, *optimized*, and the numerical results of Ref. [32], *numerical*. The analytical fit done to the numerical values is also plotted, *fitted*.

fulfill. We have included them in our functional and a

constrained search of the optimum parameters is carried out. As a result for the minimum both of them hold within 10^{-6} hartree and the minimum energy is not substantially different to that obtained in an unconstrained minimization. The tail of the effective potential is expected to be better reproduced by imposing the homo-condition.

The method has been generalized to work with multi configuration wave functions as in the Multi Configuration Hartree Fock method. This Multi Configuration OEP method provides results very close to those obtained by using Multi Configuration Hartree Fock (MCHF).

An analysis on the convergence on the parameterization of the effective potential is carried out. The results obtained are very close to the Hartree-Fock self consistent energies, eigenvalues and exchange energies improving previous optimized effective potential calculations. Results for several bound states of the beryllium atom by using multi configuration wave functions have been reported.

Acknowledgments

This work has been partially supported by the Ministerio de Ciencia y Tecnología and FEDER under contract BFM2002-00200, and by the Junta de Andalucía.

-
- [1] C. Froese-Fischer, *The MCHF method in electronic structure* (Institute of Physics Publishing, Bristol, 1997).
 - [2] R. T. Sharp and G. K. Horton, Phys. Rev. **90**, 317 (1953).
 - [3] J. C. Slater, Phys. Rev. **81**, 385 (1951).
 - [4] J. D. Talman and W. F. Shadwick, Phys. Rev. A **14**, 36 (1976).
 - [5] Y. Li, J. B. Krieger, J. A. Chevary, and S. H. Vosko, Phys. Rev. A **43**, 5121 (1991).
 - [6] E. Engel, J. A. Chevary, L. D. Macdonald, and S. H. Vosko, Z. Phys. D **23**, 7 (1992).
 - [7] S. Hirata, S. Ivanov, I. Grabowski, and R. J. Bartlett, J. Chem. Phys. **116**, 6468 (2002).
 - [8] W. Yang and Q. Wu, Phys. Rev. Lett. **89**, 143002 (2002).
 - [9] J. B. Krieger, Y. Li, and G. J. Iafrate, Phys. Rev. A **45**, 101 (1992).
 - [10] T. Grabo, T. Kreibich, S. Kirth, and E. K. U. Gross, in *Strong coulomb correlations in electronic structure calculations: Beyond the local density approximation*, edited by V. I. Anisimov (Gordon and Breach, Newark NJ, 2000), pp. 203–311.
 - [11] Y. Wang, J. P. Perdew, J. A. Chevary, L. D. Macdonald, and S. H. Vosko, Phys. Rev. A **41**, 78 (1990).
 - [12] E. Engel and S. H. Vosko, Phys. Rev. A **47**, 2800 (1993).
 - [13] J. B. Krieger, Y. Li, and G. J. Iafrate, Phys. Rev. A **46**, 5453 (1992).
 - [14] S. Kümmel and J. P. Perdew, Phys. Rev. Lett. **90**, 043004 (2003).
 - [15] P. P. Szydlík and E. S. Green, Phys. Rev. A **9**, 1885 (1974).
 - [16] J. P. Daudey and M. Berrondo, Int. J. Quantum Chem. **19**, 907 (1981).
 - [17] F. J. Rogers, B. G. Wilson, and C. A. Iglesias, Phys. Rev. A **38**, 5007 (1988).
 - [18] A. Görling and M. Levy, Phys. Rev. A **50**, 196 (1994).
 - [19] L. Fritsche and J. Yuan, Phys. Rev. A **57**, 3425 (1998).
 - [20] A. Görling, Phys. Rev. Lett. **83**, 5459 (1999).
 - [21] S. Ivanov, S. Hirata, and R. J. Bartlett, Phys. Rev. Lett. **83**, 5455 (1999).
 - [22] R. Colle and R. K. Nesbet, J. Phys. B: At. Mol. Opt. Phys. **43**, 2475 (2001).
 - [23] E. N. Lassette, J. Chem. Phys. **83**, 1709 (1985).
 - [24] X. Y. Pan and V. Sahni, Phys. Rev. A **67**, 012501 (2003).
 - [25] E. Clementi and C. Roetti, At. Data Nucl. Data Tables **14**, 177 (1974).
 - [26] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes in FORTRAN: The Art of Scientific Computing* (Cambridge University Press, Cambridge, 1992).
 - [27] J. B. Krieger, Y. Li, and G. J. Iafrate, Phys. Lett. A **148**, 470 (1990).
 - [28] S. K. Ghosh and R. G. Parr, J. Chem. Phys. **82**, 3307 (1985).
 - [29] M. Levy and J. P. Perdew, Phys. Rev. A **32**, 2010 (1985).
 - [30] F. J. Gálvez, E. Buendía, and A. Sarsa, J. Chem. Phys. **117**, 6071 (2002).
 - [31] F. J. Gálvez, E. Buendía, and A. Sarsa, J. Chem. Phys. **118**, 6858 (2003).
 - [32] K. Aashamar, T. M. Luke, and J. D. Talman, At. Data and Nucl. Data Tables **22**, 443 (1978).
 - [33] C. F. Bunge, J. A. Barrientos, and A. V. Bunge, At. Data

- and Nucl. Data Tables **53**, 113 (1993). (1991).
- [34] C. Froese-Fischer, Comput. Phys. Commun. **64**, 369